The enigmatic mossy cell of the dentate gyrus

Helen E. Scharfman^{1,2}

Abstract | Mossy cells comprise a large fraction of the cells in the hippocampal dentate gyrus, suggesting that their function in this region is important. They are vulnerable to ischaemia, traumatic brain injury and seizures, and their loss could contribute to dentate gyrus dysfunction in such conditions. Mossy cell function has been unclear because these cells innervate both glutamatergic and GABAergic neurons within the dentate gyrus, contributing to a complex circuitry. It has also been difficult to directly and selectively manipulate mossy cells to study their function. In light of the new data generated using methods to preferentially eliminate or activate mossy cells in mice, it is timely to ask whether mossy cells have become any less enigmatic than they were in the past.

Golgi technique

A method established by Camillo Golgi that stains many neurons almost completely (except for their axons) so they can be visualized in detail.

Optogenetics

The use of light to activate opsins (located in the plasma membrane), which open channels for cations or anions to flow. After opsins are expressed in one cell type, they can be activated selectively by light. Targeting opsins to specific cell types is done after identifying unique genes in the cell type, so the combination of light (opto-) and genetics (optogenetics) is fundamental to the approach.

¹Departments of Child and Adolescent Psychiatry, Physiology and Neuroscience, and Psychiatry, New York University Langone Medical Center, New York 10016, USA. ²Center for Dementia Research, The Nathan Kline Institute for Psychiatric Research, Orangeburg, New York 10962, USA. Helen.Scharfman@nki. rfmh.org

doi:<u>10.1038/nrn.2016.87</u> Published online 28 Jul 2016 The mammalian hippocampus is typically divided into areas CA1, CA2 and CA3, and the dentate gyrus. As in most cortical circuits, hippocampal neurons are classified as either glutamatergic principal cells or GABAergic interneurons. The dentate gyrus contains a primary glutamatergic principal cell type, the granule cells, and is distinct from other hippocampal regions in having an additional glutamatergic principal cell type, the mossy cells. Mossy cells are named for their characteristic 'mossy' appearance when stained using the Golgi technique, which reveals clusters of complex spines, known as thorny excrescences, on their proximal dendrites. Mossy cells comprise a large subset of neurons of the dentate gyrus and are implicated in several pathological conditions, but these cells have been hard to define functionally. Many hypotheses about their functions have been proposed, but mossy cells have remained 'enigmatic' because of the persistent uncertainty about their functions.

Recent technical advances that enable the selective manipulation of mossy cells^{1,2} mean that this enigmatic past is giving way to a clearer understanding. For example, it is now evident that mossy cells can both excite granule cells and indirectly inhibit them by activating GABAergic interneurons. However, the new results raise more questions than they seem to answer. Therefore, it is timely to consider the new data in the context of the pre-existing hypotheses about mossy cells. In this Review, I first provide an overview of the dentate gyrus circuitry and describe the existing hypotheses about mossy cell function. I then explain how the new data using transgenic mice and optogenetics have shed light on these hypotheses, and I suggest additional experiments that are necessary to resolve the outstanding questions.

The dentate gyrus

Structure, circuitry and cell types. The basic structure of the dentate gyrus is discussed in detail elsewhere³ and summarized in FIG. 1a,b. The dentate gyrus is composed primarily of granule cells, which are oriented in a stereotypical manner. The dendrites of granule cells are present in the molecular layer, and their cell bodies form the adjacent granule cell layer (GCL). Between the GCL and area CA3, there is a polymorphic layer called the hilus (FIG. 1) that contains the granule cell axons, which are called mossy fibres (FIG. 1c). The molecular layer is divided into the outer molecular layer (OML), the middle molecular layer (MML) and the inner molecular layer (IML). This anatomical organization is similar in rodents and primates (FIG. 1a,b).

Afferents to the dentate gyrus come from many sources (FIG. 1c; TABLE 1). The primary cortical input to the dentate gyrus is the glutamatergic projection from layer II of the entorhinal cortex (also known as the perforant path), which is responsible for most of the afferent inputs to the OML and MML³⁻⁵. Mossy cell axons are a major afferent input to the IML⁶⁻⁸. The outputs of the dentate gyrus are from the granule cell mossy fibres that project to area CA3 (REF. 3) (FIG. 1c; TABLE 1).

The major cell types of the dentate gyrus include the granule cells, mossy cells and GABAergic interneurons³ (FIG. 2). Most granule cells are located in the GCL, but there are small subsets in the IML (known as semilunar granule cells^{9,10}) and hilus (known as ectopic granule cells¹¹) (FIG. 2a). Stem cells are located in the subgranular zone¹². These subgranular zone progenitors divide throughout life and migrate primarily to the GCL where they become granule cells and integrate into the dentate gyrus circuitry in a similar manner to granule cells born in early life¹².



Figure 1 | The organization of the dentate gyrus of rodents and primates. a | The general organization of the dentate gyrus in the coronal and horizontal planes of the rodent brain. The area surrounded by the dashed box is expanded in the inset figure to show the laminar organization of the dentate gyrus, which is composed of a molecular layer (MOL), granule cell layer (GCL) and hilus (HIL). The GCL contains the principal cells of the dentate gyrus, the granule cells. The dendrites of the granule cells extend into the MOL, and their axons traverse the HIL and terminate in area CA3c. **b** | A schematic of the primate dentate gyrus shows that it is similar to that of the rodents, but that there is gyrification. In addition, CA3c is larger in primates than in rodents and includes a reflected blade (indicated by the arrowhead). c | The layers of the dentate gyrus are shown. The sources of major afferent inputs are shown on the right (red box indicates the GABAergic input and grey boxes indicate inputs from other neurotransmitters). GABAergic interneurons innervate all layers. The lateral entorhinal cortex (LEC) and the medial entorhinal cortex (MEC) innervate the outer molecular layer (OML) and the middle molecular layer (MML), respectively.

Supramammillary (Mamm), cholinergic, mossy cell and ventral CA3 (vCA3) pyramidal cell axons innervate the inner molecular layer (IML). The OML and MML also receive inputs from the brainstem (including noradrenergic and 5-hydroxytryptamine (5-HT; also known as serotonin) inputs) and from basal forebrain cholinergic neurons. The HIL receives diverse inputs, including the axons of granule cells, dentate gyrus GABAergic neurons, mossy cells, CA3 pyramidal cells, neuromodulatory inputs from the brainstem (such as noradrenaline (NA), 5-HT and dopamine (DA)) and basal forebrain cholinergic neurons^{76,91,92,132,133}. The efferents from the dentate gyrus to other areas arise mainly from granule cells that project to the HIL and CA3. The granule cell axon, called a mossy fibre, is complex. It makes giant boutons that innervate thorny excrescences of mossy cells and pyramidal cells and small boutons that arise from hilar collaterals and filamentous extensions from the giant boutons^{25,29,30,33,34}. The small boutons primarily contact interneurons but also form contacts on distal dendrites of mossy cells^{7,34}. ACh, acetylcholine; F, fissure; SGZ, subgranular zone.

The interneurons in the dentate gyrus are diverse, with similarities to the interneurons in other hippocampal subfields¹³ (FIG. 2b). They are commonly classified by the location of their cell body and axon projection, a classification scheme that emphasizes the specificity of many interneuron terminal fields for a sublayer of the dentate gyrus^{13–15}. This is similar to the way in which the interneurons in areas CA1–CA3 are classified¹³, but the nomenclature is distinct^{13–15} (FIG. 2b). For example, a major population of hilar neurons have a hilar cell body and project to the outer two-thirds of the molecular layer. They are called HIPP cells (hilar cells that project to the terminal zone of the perforant path^{13–15}) (FIG. 2b). However, the dentate gyrus interneuron axons do not always terminate precisely in one layer^{16,17}, particularly during development, before pruning of dendrites and axons¹⁸. In adulthood, the lack of specificity of some axons may be important for the maintenance

Table 1 Afferents and efferents of the dentate gyrus								
Hippocampal afferents		Extra-hippocampal	Efferents to dentate gyrus		Efferents			
Glutamatergic	GABAergic	afferents	Glutamate	GABA	to CA3			
Outer molecular layer								
Not applicable	 HIPP¹³⁻¹⁵ MOPP¹³⁻¹⁵ NG cell¹³⁵ 	• LPP ^{4,5} • NA ⁹¹ • 5-HT ⁹¹	Not applicable	• MOPP ¹³⁻¹⁵ • NG cell ¹³⁵	Not applicable			
Middle molecular layer								
Not applicable	 HIPP¹³⁻¹⁵ MOPP¹³⁻¹⁵ NG¹³⁵ 	 MPP^{4,5} NA⁹¹ 5-HT⁹¹ 	Not applicable	• MOPP ¹³⁻¹⁵ • NG ¹²³	Not applicable			
Inner molecular layer								
• MC ^{6-8,25,26,49,41} • vCA3 (REF. 81)	• HICAP ^{13,15}	 ACh¹³⁸ Mamm^{139–141} 	• smGC ^{9–10}	Multiple ¹⁶	• smGC ^{9–10}			
Granule cell layer								
Not applicable	• BC ¹³⁻¹⁵ • AA cell ¹³⁻¹⁵	• Mamm ¹³⁹	• mGC ^{3,31} • abGC ¹²	• BC ¹³⁻¹⁵ • AA cell ¹²⁶	• mGC ^{3,31} • abGC ¹²			
Hilus								
• GC ^{3,34} • MC ^{7,8,35} • CA3 (REFS 74–76,78)	• Multiple ¹⁶	 NA^{78,91,92} 5-HT⁹¹ DA^{91,92,132} ACh¹³³ 	 MC^{7,8,35} eGC¹¹ smGC⁹⁻¹⁰ mGC^{3,31} abGC¹⁴⁰ 	 HIPP¹³⁻¹⁵ HICAP¹³⁻¹⁵ Other¹⁶ 	• eGC ¹¹			

The table lists the neuronal subtypes that innervate each layer of the dentate gyrus. 5-HT, 5-hydroxytryptamine; AA, axo-axonic; abGC, adult-born granule cell; ACh, acetylcholine; BC, basket cell; DA, dopamine; eGC, ectopic granule cell; GC, granule cell; HICAP, interneurons with a hilar cell body with axon targeting the commissural/associational pathway; HIPP, hilar cells that project to the terminal zone of the perforant path; LPP, lateral perforant path; Mamm, supramammillary; MC, mossy cell; mGC, mature granule cell; MOPP, interneurons with a molecular layer cell body that project to the terminal zone of the perforant path; MPP, medial perforant path; NA, noradrenaline; NC, neurogliaform; smGC, semilunar granule cell; vCA3, ventral CA3.

of inhibition in the event that a subset of interneurons is damaged. Indeed, some interneuron subtypes are more vulnerable to insults and injury than others, and the dentate gyrus interneurons can also be classified according to their relative resistance to injury. A common example of a relatively resistant cell type is the parvalbumin (PV)-expressing interneuron, which has an axon that forms a plexus around the granule cell somata. These interneurons are often called basket cells, or are included in the larger group of perisomatic-targeting neurons. Basket cells in the dentate gyrus are considered to be resistant to seizures because they survive the excitotoxicity that kills other cell types of the dentate gyrus, such as the HIPP cells¹⁹⁻²¹. The dentate gyrus interneurons have also been classified according to their expression of calcium-binding proteins (such as PV or calbindin²⁰) or neuropeptides (including somatostatin, neuropeptide Y and cholecystokinin²²). Although most dentate gyrus interneurons project locally (in the vicinity of their cell body), basket cells and some neuropeptide Y-expressing cells project to distant sites, such as the contralateral dentate gyrus^{23,24}.

Mossy cells are located only in the hilus^{3,7,25}. Their dendrites usually span the hilus and sometimes extend into the molecular layer^{6,7,25–28} (FIG. 2a). One hallmark of mossy cells is the presence of thorny excressences, which do not exist on interneurons or granule cells^{6,25,26}. These large spine complexes are postsynaptic to extremely large synaptic boutons of granule cells that are densely packed

with glutamatergic vesicles (the so-called massive or giant mossy fibre boutons²⁹⁻³²). The term 'mossy', used for both mossy cells and granule cell mossy fibres, can be confusing but is appropriate, given the 'mossy' appearance of both. The general arrangement of the granule cell boutons and the thorny excrescences is similar in the hilus and CA3: massive boutons are opposed to thorny excrescences on mossy cells or pyramidal cells, whereas smaller boutons of granule cells are located on many hilar collaterals of mossy fibres, where they contact hilar interneurons and distal dendrites of mossy cells⁷. Small boutons are also found on filamentous extensions from giant boutons^{33,34}.

Many investigators assume that thorny excrescences define mossy cells, but there are spiny hilar cells without thorns that have the same physiological characteristics as 'thorny' mossy cells^{7,35,36}. Indeed, mossy cells vary in the degree to which they have thorny excrescences both within a given species and across species^{7,28,35,37}. They also vary in their expression of neurochemical markers; for example, calretinin is expressed in ventral but not dorsal mossy cells in mice^{38,39}. Thus, mossy cells are best defined using more criteria than only the presence of thorny excrescences⁷ (TABLE 2).

The mossy cell axon projection is complex (FIG. 3). A large projection (known as a 'distant' or 'intralamellar' projection) terminates away from the cell body in both the ipsilateral and contralateral dentate gyrus^{6–8,26,40,41}.

HIPP

10 µm





HIL



75 µm

Figure 2 | The cell types of the dentate gyrus. a | Glutamatergic cells of the dentate gyrus include granule cells and mossy cells. Granule cells are not only located in the granule cell layer (GCL); there are small subsets in the inner molecular layer (IML) and hilus (HIL), and precursors to granule cells are located in the subgranular zone (SGZ). Mossy cells have long dendrites, some of which extend into the molecular layer (MOL; comprised of the IML, middle molecular layer (MML) and outer molecular layer (OML))^{27,28}. b | GABAergic neurons of the dentate gyrus are heterogeneous. Their nomenclature is based on the location of the cell body and the axon terminal field¹³⁻¹⁵. For example, MOPP cells have a cell body in the MOL and terminals in the OML and MML, where the terminals of the perforant path are located¹³⁻¹⁵. HICAP cells (interneurons with a hilar cell body with axon targeting the commissural/associational pathway) innervate the IML, where the commissural/associational projection from mossy cells is located¹³⁻¹⁵. The neurons that innervate the granule cell somata or the axon initial segments are called perisomatic-targeting cells. Two of the most common cell types in this group are basket cells, which make basket-like endings around the granule cells and often have a pyramidal-shaped soma that is located at the border of the GCL and HIL¹³⁴, and axo-axonic (AA) cells¹³⁻¹⁵. AA cells are often present near or in the GCL, as shown, and innervate granule cell axon initial segments¹³⁻¹⁵. Several GABAergic neuron subtypes innervate granule cell dendrites. The most common of these are cells in the HIL that innervate the OML and MML (HIPP cells, hilar cells that project to the terminal zone of the perforant path). An example of a neurogliaform cell (NG) that innervates the molecular layer is shown¹³⁵. There are some types of GABAergic neurons that have an axon that innervates more than one layer (here labelled 'other')^{16,17}, and some interneurons innervate each other, such as calretinin-expressing hilar cells (CR)¹³⁶. c | The upper image shows an example of a biocytin-filled mossy cell with thorny excrescences labelled in the higher magnification inset (indicated by the arrows). The lower image shows an interneuron that was filled with biocytin in a rat hippocampal slice. Arrows point to primary dendrites that are smooth relative to the mossy cell. F, fissure.

Table 2 Characteristics of mossy cells									
General characteristic	Specific characteristic	Granule cell	Mossy cell	Dentate gyrus interneuron	Refs				
Anatomical characteristics									
Cell body	Shape	Oval or circle	Diverse	Diverse	3,6-8,25,26				
	Size	Small	Usually large	Usually small					
Axon projections and targets	Local	Hilus and CA3	Hilus	All layers, usually subtype specific	3,6–8,25,26				
	Distal	None	Inner molecular layer	Weak					
	Contralateral	None	Inner molecular layer	Weak (BCs and HIPP cells)					
Dendrites	Location	Molecular layer	Mostly hilar	Diverse	3,6-8,25,26				
	Spines	Dense	Dense, usually thorny	Usually few					
Immunohistochemistry	GluR2 or GluR3	Yes	Yes	No	96,142				
	Calretinin	Yes, in young GCs	Yes, in mice	Yes, but not in all interneurons	38,39,70,141,143				
	CGRP	No	Yes	No	144				
	CART	No	Yes, in human	No	37				
	D2R	No	Yes	Yes, in a rare subtype of interneurons	59				
	CB1R	No	Yes	Yes	67				
	p11	No	Yes	Yes, in BCs	119				
	Dysbindin 1C	No	Yes	Unclear	120				
	Glucocorticoid type 2 receptor	Yes	Yes	No	145				
	Calbindin	Yes	No	Yes, in a subtype of interneurons	13,20,126				
	Parvalbumin	No	No	Yes, in BCs	13,20,126				
	STEP	No	Yes	Yes, in HIPP cells	128				
Electrophysiological characteristics									
Intrinsic	RS, FS or SS?	RS	RS	FS or SS	7,13,16,17,146,147				
	Action potential dv/dt	>1	>1	~1					
	AHP	Small	Small	Large					
	Time constant	Short	Long	Short					
	Sag	No	Some	Some					
	SFA	Robust	Some	Weak					
Synaptic	Spontaneous EPSCs	Infrequent, relative to hilar cells	Frequent, large	Variable, usually small and fast	35,36,55,148,149				
	Response to trains of >2 Hz PP stimuli	One response occurs per stimulus	Depolarization of the cell occurs during the train and outlasts the train	Variable	112,150,151				
	Response to ACh or DA	Hyperpolarization	Depolarization	Variable (unclear for DA)	94,132,137				
Other characteristics									
Developmental origin		Similar to MCs	Similar to GCs	Distinct from MCs and GCs	152				
Relative vulnerability		Resistant	Vulnerable	Resistant (BCs) or sensitive (HIPP cells)	7,20,107,126				

ACh, acetylcholine; AHP, afterhyperpolarization; BCs, basket cells; CART, cocaine-amphetamine regulating transcript; CB1R; cannabinoid type 1 receptor; CGRP, calcitonin gene-related peptide; DA, dopamine; dv/dt, rate of rise (ratio of decay of an action potential at threshold for spike generation); D2R, dopamine D2 receptor; EPSCs, excitatory postsynaptic currents; FS, fast spiking; GCs, granule cells; GluR2, glutamate receptor type 2; HIPP, hilar cells that project to the terminal zone of the PP; MC, mossy cell; PP, perforant path; RS, regular spiking; Sag, a reduction in a voltage response to a persistent current command; SFA, spike frequency adaptation; SS, slow spiking; STEP, striatum-enriched protein-tyrosine phosphatase.

Electron microscopy

The use of microscopes with very high (nanometre) resolution, made possible by accelerating the electrons through a specialized microscope. Electron microscopy can be used with very thin brain sections, allowing parts of neurons (such as synapses) to be detected.

Contextual fear conditioning A behavioural test that

examines the response to an environment or context after a prior exposure to the context and a painful stimulus. The terminals of this distant projection primarily synapse in the IML, and electron microscopy shows that they primarily innervate spines⁸. Because interneurons with dendrites in the IML rarely have spines, it is likely that the distant mossy cell axon projection primarily innervates granule cells⁸. Mossy cell axons also collateralize in the hilus and have extensions in the MML and OML^{7,8,35}. Near the cell body, the hilar collaterals are robust, and there is also innervation of the IML^{35,42}. In summary, a mossy cell innervates more than one dentate gyrus layer as well as both local and distant parts of the dentate gyrus, making it hard to predict its functions *in vivo*.

The role of the dentate gyrus in behaviour. To date, the dentate gyrus has been implicated in several important behavioural functions. These functions have been segregated into those that are often called 'cognitive', which are dependent on the more dorsal or septal half of the dentate gyrus, and those that are more 'emotional', which depend on the ventral or temporal half of the dentate gyrus⁴³. The cognitive functions include those related to spatial memory, whereas the emotional functions include those associated with the regulation of mood44 and anxiety or with behaviours that have a component of stress or fear (such as contextual fear conditioning⁴⁵). Additional functions have also been attributed to the dentate gyrus^{44,46}. These divisions assume that neurons of the dorsal dentate gyrus are the most important for cognitive functions; however, it is important to note that ventral mossy cells may also contribute to these functions through their major afferent inputs to the dorsal IML (FIG. 3). Indeed, because the IML is proximal to the granule cell somata, mossy cell terminals in this region are in an ideal position to influence the activity of granule cells.

A function of the dentate gyrus that has been discussed a great deal is pattern separation, which is the ability of the dentate gyrus network to receive a pattern of afferent inputs and 'separate' them so that the outputs are less similar than the inputs^{46–50}. It has been suggested that pattern separation in the dentate gyrus is important for memory storage in area CA3 because it allows similar experiences to be stored in different subsets of CA3 pyramidal cells, thus facilitating accurate memory retrieval⁴⁸. This theory assumes that the afferent input patterns that are subject to pattern separation arrive at the dentate gyrus via the perforant path. However, mossy cells provide a major glutamatergic input to granule cells (see below), suggesting that they could send input patterns to granule cells in addition to, or instead of, the perforant path.

Mossy cell function in the dentate gyrus

Do mossy cells excite or inhibit granule cells? Experiments to address the question of whether mossy cells are excitatory or inhibitory started in the 1980s^{51,52}; however, it was not until the mid-1990s that conclusive evidence was provided that mossy cells are glutamatergic⁵³ and can excite granule cells through direct inputs⁵⁴. The net effect of mossy cell input on granule cells remains unclear because mossy cells also activate dentate gyrus interneurons that inhibit granule cells^{54,55}.

One problem faced by researchers attempting to address this question is that it is difficult to selectively activate mossy cells using stimulating electrodes: mossy cell axons and dendrites are spatially close to other cells of the dentate gyrus and to axonal projections that influence the dentate gyrus. One strategy to circumvent this problem has been to record from pairs of monosynaptically connected mossy cells and granule cells⁵⁴. This approach



Figure 3 | **Organization of the mossy cell axon. a** | The axon of a single ventral mossy cell is illustrated schematically. Near the soma, the local ipsilateral branches of the mossy cell axon make synapses in the hilus (HIL) and the inner molecular layer (IML)^{7,35,42}. Far from the soma, the distant ipsilateral or distant contralateral branches of the axon project primarily to the IML^{6-8,25,40,41}. **b** | Distant and local ipsilateral circuitry. At distant ipsilateral locations, the axon primarily makes synapses on spines; because there are few spines on dentate gyrus interneurons, it is likely that the mossy cell axon in the IML innervates granule cells. In addition, the mossy cell axon extends hilar,

outer molecular layer (OML) and middle molecular layer (MML) collaterals^{7,8,35}. It is not clear whether mossy cell terminals in the OML and MML make synapses, or what cell types mossy cells target in the HIL (indicated by question marks). At local ipsilateral locations, the mossy cell axons collateralize in the HIL, especially near the soma. There, they are likely to contact interneurons because of the numerous interneuron dendrites present in this region and the absence of granule cell dendrites. Mossy cells also make local projections to the IML, but these are not as numerous as those to distant sites⁸. GCL, granule cell layer; SGZ, subgranular zone.

Cre recombinase

(Cre). Part of a site-specific recombination system derived from *Escherichia coli* bacteriophage P1. Two short DNA sequences (*loxP* sites) are engineered to flank the target DNA. Activation of the Cre recombinase enzyme catalyses the recombination between the *loxP* sites, leading to excision of the intervening sequence.

Disinhibition

A decrease in inhibition (usually GABAergic). For example, blockade of the release of GABA from GABAergic neurons would result in a decrease in inhibition of the neuron that is postsynaptic to the GABAergic neuron.

Long-term potentiation

(LTP). A lasting increase in synaptic transmission. LTP is often elicited by a brief period of high-frequency presynaptic firing. However, other types of stimulation can elicit LTP, such as exposure to neuromodulators.

Field potentials

The changes in the extracellular potential that reflect changes in the flow of cations and anions in the extracellular space.

Retrograde signalling

The changes induced in a presynaptic terminal, usually mediated by a neuromodulator acting on its presynaptic receptors, which are evoked by release of the neuromodulator from the postsynaptic site. showed that the monosynaptic mossy cell input to a granule cell generated an excitatory postsynaptic potential (EPSP) that could only be detected when GABAergic inhibition was blocked. The mossy cell input seemed to be weakly excitatory because failures of synaptic transmission were frequent⁵⁴. This suggested that the primary effect of mossy cells is the inhibition of granule cells. However, when the postsynaptic granule cell was depolarized, the effects of presynaptic mossy cell input were larger; thus, mossy cells can have a robust excitatory effect on depolarized granule cells⁵⁴. Additional studies using approaches that lesioned mossy cells or the molecular layer in hippocampal slices^{56,57,58} provided further evidence that mossy cells excite granule cells.

Despite the value of these findings, it has become increasingly important to find a way to selectively interrogate mossy cells. Therefore, the emergence of two transgenic mouse lines that express Cre recombinase (Cre) relatively specifically in mossy cells has been an important advance. In one of these mouse lines, the Crlr promoter (Crlr encodes calcitonin receptor-like receptor) was used to drive Cre expression¹. Although this approach produced selective Cre expression in mossy cells within the dentate gyrus, CA3 neurons also showed some Cre expression¹. In the second mouse line, the Drd2 promoter (Drd2 encodes the dopamine D2 receptor) was used to drive Cre expression. These mice also showed strong Cre expression in mossy cells; however, an additional population of hippocampal interneurons also expressed the Drd2 promoter, showing that the Cre expression in mossy cells was not selective^{59,60}.

To date, mossy cell function has only been addressed using the Crlr-Cre mouse line¹. These mice were crossed with mice carrying a floxed diphtheria toxin receptor, and diphtheria toxin was administered in adult mice to kill mossy cells. One week after toxin injection, mossy cells were deleted and granule cells showed disinhibition in response to a perforant path stimulus, supporting the hypothesis that the effect of mossy cells is primarily inhibitory to granule cells. However, both spontaneous excitatory postsynaptic currents (EPSCs) and inhibitory postsynaptic currents (IPSCs) were decreased in patched granule cells in brain slices from these mice. Furthermore, the disinhibitory effects on granule cells driven by the deletion of mossy cells did not persist; the authors suggested that this was a result of compensatory changes^{1,61}.

The problem of potential compensatory changes in response to cell death can now be circumvented by activating mossy cells selectively with optogenetics. To date, one study has been published in which the excitatory opsin channelrhodopsin was expressed in commissurally projecting dentate gyrus neurons², which are mainly mossy cells. The authors showed that both EPSCs and IPSCs were evoked in granule cells in response to single light pulses focused on the commissurally projecting terminals, suggesting that mossy cells had both excitatory and inhibitory effects on granule cells. However, when the light stimulus was timed to precede perforant path stimulation, the response of the granule cells to the perforant path input was reduced. This suggested that

the dominant effect of mossy cells on perforant pathevoked responses of granule cells is inhibitory. Repetitive light pulses that were more similar to patterns of activity observed in vivo were also tested and revealed that the excitation of GABAergic neurons was more persistent than the excitation of granule cells. Thus, the inhibitory effects of mossy cells might be stronger than their excitatory effects in vivo. In addition, the results showed that mossy cells are likely to activate a range of interneuron cell types, with a preference for basket cells. These experiments provided the most compelling evidence to date that the primary effect of mossy cells is the inhibition of granule cells. However, it is important to note that the commissurally projecting neurons of the dentate gyrus targeted in this study also include inhibitory neurons^{23,24}. In addition, the studies were conducted in hippocampal slices, not in vivo.

Adult neurogenesis. Very few studies have examined the influence of mossy cells on adult neurogenesis. It has been suggested that mossy cells could be important for adult neurogenesis because they provide the first gluta-matergic input to adult-born granule cells⁶². However, the deletion of mossy cells in *Crlr*–Cre mice treated with the diphtheria toxin did not appear to have a major effect on adult-born neurons¹ (although many characteristics of the newborn neurons were not tested).

Contribution to plasticity. Additional insight into the potential role of mossy cells has come from studies showing that mossy cell synapses exhibit long-term potentiation (LTP). For example, granule cell input to mossy cells exhibits LTP with characteristics similar to those of mossy fibre-evoked LTP in CA3 pyramidal cells63. However, it is unclear whether mossy cell input to granule cells also shows LTP. One in vivo study showed LTP of field potentials recorded in the IML after high-frequency stimulation of the hilus64. However, another in vivo study using IML stimulation did not find LTP of the mossy cellgranule cell pathway65. Notably, high-frequency stimulation of the perforant path evoked both LTP of the perforant path-granule cell synapse and heterosynaptic LTP of the IML-evoked response, suggesting that the mossy cell input to granule cells can be potentiated⁶⁵. Another in vivo study showed that LTP of the perforant path-granule cell synapse leads to LTP of the contralateral mossy cell-granule cell synapse⁶⁶. This study was important because it showed that LTP of the perforant pathgranule cell synapse induces plasticity of 'downstream' mossy cells and thus appears to be permissive for LTP of the mossy cell-granule cell synapse.

A potential explanation for the lack of LTP at the mossy cell–granule cell synapse that was observed in the earlier study⁶⁵ was provided by experiments showing that the depolarization of granule cells suppresses the mossy cell input to granule cells, an effect that was mediated by the retrograde signalling of endocannabinoids^{67,68} through cannabinoid receptors (TABLE 2). Thus, the mossy cell input to granule cells can be suppressed by high-frequency stimulation that sufficiently depolarizes the granule cell.

Other studies of mossy cell synaptic plasticity suggested that mossy cells influence the LTP of the perforant path-granule cell synapse. For example, one study⁶⁹ took advantage of the fact that in mice calretinin is expressed primarily in mossy cells^{38,39}. LTP of the perforant path-granule cell synapse was reduced in mice lacking calretinin compared with wild-type mice⁶⁹. Although intriguing, the study was limited by the constitutive nature of the knockout and the expression of calretinin in cells other than mossy cells (including a subtype of interneuron in the dentate gyrus⁷⁰). However, other methods have also shown that mossy cells may facilitate or even be required for LTP of the perforant path input to granule cells. Voltage imaging in hippocampal slices showed that the granule cell-mossy cell-granule cell circuit was necessary for perforant path-granule cell LTP58. Furthermore, it was shown that the expression of growth-associated protein 43 (GAP43; also known as neuromodulin) was increased in mossy cells in response to high-frequency stimulation of the perforant path, which could support the persistence of LTP⁷¹. Thus, it has been suggested that the mossy cell and perforant pathway inputs to granule cells are cooperative65,71.

Role in hippocampal oscillations. Hippocampal neurons display oscillatory behaviour at various frequencies in vivo, and these rhythms are thought to have a role in the encoding of spatial information⁷². Therefore, one way to better understand the role of mossy cells in the hippocampal network is to determine how variations in their activity relate to those recorded simultaneously in the hippocampal electroencephalogram (EEG). Two studies have shown that, in anaesthetized animals, mossy cells discharge in a similar manner to principal cells in relation to theta oscillations73,74. Both studies pointed out that mossy cells might act to phaselock the activity of granule cells to the EEG throughout the hippocampal dorsoventral axis⁷³; this is consistent with the idea that mossy cells link different subsets of granule cells⁷⁵ (FIG. 4a).

Another study⁷⁴ recorded the intracellular activity of mossy cells in anaesthetized rats. The authors found that when the tail was pinched during ongoing theta oscillations, mossy cells could, but did not always, depolarize. This variability is consistent with the idea that there are functional subtypes of mossy cells, which is also suggested by variations in their structure⁷ and by dorsoventral variations in the expression of calretinin (see above)^{38,39}.

In this context, it is interesting to consider the data from *Crlr*–Cre mice treated with the diphtheria toxin¹ that showed that, shortly after treatment with the toxin, theta power in the dentate gyrus increased. This implies that the activity of the dentate gyrus interneurons was suppressed following the loss of mossy cells, because other methods that depress the firing of the dentate gyrus interneurons also increase theta power⁷⁶. These data suggest that mossy cells are likely to have a more important role in theta rhythm than previously considered.

Possible contributions to behaviour

Although our understanding of the role of mossy cells in dentate gyrus function is limited, several intriguing hypotheses about their potential contributions to behaviour have been proposed, based on the knowledge that we do have.

Associative functions. One early hypothesis for the role of mossy cells in hippocampal function is reflected in one of the names given to their axon projection, the associational projection⁷⁵. Mossy cells may allow physically separate subsets of granule cells to be associated with one another, analogous to the way in which the recurrent collaterals of CA3 pyramidal cells allow different subsets of pyramidal cells to interact⁷⁵ (FIG. 4a). This idea was supported by detailed electron microscopy-based studies of the mossy cell axon^{8,42}, which showed that, because of the numerous varicosities present on a mossy cell axon in the IML, a single mossy cell could potentially link multiple granule cells⁷⁵.

Pattern separation. As described above, a major question is how the dentate gyrus separates patterns of afferent input that contain elements that are identical (overlapping). An example of such overlap would be two afferent inputs that are active during two different patterns of input⁵⁰; however, other characteristics could potentially also be shared. Mossy cells may contribute to the separation of two overlapping patterns in several ways. For two patterns of input from the entorhinal cortex, separation could occur if the shared (identical) element of one pattern was coincident with mossy cell input to granule cells but not coincident with the other pattern. When coincident, mossy cells could depolarize the same granule cell dendrite as the entorhinal input and do so at a similar time, causing action potentials to be elicited by one pattern of input but not the other. Alternatively, activity of mossy cells in one pattern but not the other pattern could increase GABAergic inhibition of granule cells, reducing the number of action potentials in the granule cells in response to one pattern but not the other (FIG. 4b). One might argue that granule cells are strongly inhibited regardless of the activity of the mossy cells, because their soma and dendrites are well innervated by axons of diverse interneuron subtypes¹³⁻¹⁶. However, unlike interneurons, mossy cells could activate GABAergic interneurons broadly and synchronously, and therefore increase GABAergic inhibition of granule cells in a unique way (FIG. 4b). This idea is consistent with the proposed mossy cell-mediated phase-locking of granule cells73,74.

One argument against this idea is that the projection of mossy cells to the distant ipsilateral IML targets many more granule cells than interneurons⁸. However, the function of these excitatory inputs might differ from that of a typical excitatory input because mossy cell input to granule cells may be excitatory only under certain conditions (FIG. 4c).

Alternatively, excitatory input of mossy cells to granule cells could contribute to pattern separation by promoting the activity and development of young adultborn granule cells, and by having a weakened effect as the granule cells age. This idea is consistent with the evidence

Voltage imaging

The identification of neuronal activity by capturing changes in fluorescence that are proportional to changes in membrane potential. Typically, a voltage-sensitive dye is applied to the preparation of neurons so that voltage imaging can be conducted.

Electroencephalogram

(EEG). Recordings of the electrical activity of the brain with electrodes that are not inside the neurons (intracellular), but outside (extracellular) or remote (on the brain surface or skull).

Oscillations

The intermittent activity of neurons that is sufficiently synchronous to induce rhythmic fluctuations in the extracellular potential.

Recurrent collaterals

The branches of the axons of a population of neurons that innervate the dendrites of the same population of neurons. In area CA3, the pyramidal cells do not innervate their own dendrites but the dendrites of other CA3 pyramidal neurons.



Figure 4 | **Possible roles of mossy cells in dentate gyrus function. a** | A long-standing proposal is that mossy cells link subsets of granule cells that are spatially separated and therefore facilitate associative

learning⁷⁵. **b** | Another long-standing view is that granule cells primarily activate GABAergic interneurons, particularly basket cells^{106,112}. This could lead to a possible circuit, as shown, in which mossy cells amplify the normal feedback inhibition of granule cells by activating GABAergic interneurons. In the figure, this circuit is exemplified by the perforant path (PP)-mediated depolarization of a granule cell. If the depolarization evokes an action potential, the granule cell would subsequently depolarize a basket cell that innervates numerous granule cells (only one granule cell is depicted), increasing the inhibition of granule cells. On the right, this physiology is schematized. Without the mossy cell in the circuit (top), a PP input (indicated by the arrow) elicits an excitatory postsynaptic potential (EPSP) in the granule cell. The EPSP is followed by an inhibitory postsynaptic potential (IPSP) that reflects the feedback inhibition (that is, the activation of the basket cell by the granule cell, which in turn hyperpolarizes the granule cell). Notably, it is unclear whether basket cells innervate the same granule cells that activate them; the circuit in the diagram is simplified for clarity. With the mossy cell in the circuit (bottom), the PP input elicits an EPSP followed by a larger and longer IPSP in the granule cell because mossy cells are first depolarized by granule cells and then activate basket cells. Note that if the PP directly innervates basket cells and mossy cells, this would lead to feedforward inhibition of granule cells. c | According to another hypothesis, the

b Amplifier Granule cell response to PP input (1) Without mossy cell With mossy cell With mossy cell to PP input (1)



Granule cell response to PP input (↑)

Without mossy cell

↑ With mossy cell

excitatory effects of mossy cells depolarize granule cells only under some conditions. In this example, the effects of a CA3 pyramidal cell input to a granule cell depend on the presence or absence of concurrent cholinergic input. Without the cholinergic input, CA3 pyramidal cells will primarily hyperpolarize granule cells by activating interneurons. With cholinergic input mediated by acetylcholine (ACh), which preferentially suppresses the interneurons¹³⁷, CA3 pyramidal cells will primarily depolarize granule cells by activating mossy cells^{80,83}. The schematic includes several simplifications for clarity. For example, the cholinergic input to pyramidal cells and mossy cells is not shown. In addition, the cholinergic input depolarizes mossy cells⁹⁴, which would lead to a greater potential to excite granule cells in the circuit that is shown. d | Finally, mossy cells have also been proposed to act as 'sentinels' that inform granule cells that changes have occurred even in the absence of strong PP input. This sentinel function would be reflected by the lack of a response of a granule cell to the PP input unless the mossy cell is activated. This might occur, for example, if the PP input to granule cells is weak. Under these conditions, the PP might still activate semilunar granule cells because they are activated more readily by the PP. These cells have a robust excitatory effect on mossy cells^{9,10}, which would then activate granule cells, as shown in the diagram. On the right, this is depicted by a lack of depolarization in a granule cell (located in the granule cell layer (GCL)) in the absence of mossy cells. HIL, hilus; IML, inner molecular layer; MML, middle molecular layer; OML, outer molecular layer; SGZ, subgranular zone.

that behavioural tasks that are suggested to involve pattern separation are impaired when young adult-born granule cells are ablated, an effect that is not evident after destruction of older granule cells^{77,78}. Whether such a correlation between changes in mossy cell input and granule cell ageing occurs is unknown, but it would be consistent with the suggestion that adult-born granule cells become less active as they age⁷⁹.

Pattern completion refers to the retrieval of a stored memory when presented with a subset of the initial input that was used to produce the memory. Pattern separation and pattern completion are often discussed as if they involve the dentate gyrus and area CA3, respectively, with the granule cell mossy fibres making a one-way path to CA3. However, CA3 pyramidal cell axons also project 'back' to the dentate gyrus⁸⁰. CA3 pyramidal cell axon collaterals project far into the hilus^{81,82} and, in the ventral pole of the hippocampus, their projections extend further, into the IML⁸¹. Most of the projections arise from CA3c, with fewer arising from CA3b and CA3a^{81,82}. However, recurrent collaterals between CA3a-CA3b pyramidal cells and CA3c pyramidal cells can make it possible for CA3a-CA3b to influence the dentate gyrus via the CA3c⁸⁰. In the hilus, CA3 pyramidal cells innervate mossy cells and interneurons (most dentate gyrus interneurons have a soma or dendrites in the hilus)⁸³⁻⁸⁵. In hippocampal slices, the major effect of CA3 firing is GABAergic inhibition of granule cells⁸³. However, the delay between a pyramidal cell action potential and the onset of granule cell inhibition (10-20 ms) suggests that a simple pyramidal cell-interneuron-granule cell pathway is not responsible; instead, a pyramidal cell-mossy cell-interneuron-granule cell pathway or a pyramidal cell-pyramidal cell-interneuron-granule cell route seems more likely to be responsible⁸³. Notably, a robust pyramidal cell-mossy cell-granule cell excitatory pathway is revealed when GABAergic inhibition is blocked⁸³.

The backprojection from CA3 to the dentate gyrus might affect circuitry in the dentate gyrus and CA3 (REFS 86,87), and the existence of this pathway might add to the potential ways in which information may be processed. Although the recurrent collateral network of CA3 is often thought to provide the only robust excitatory recurrent circuitry, synapses made by the CA3–dentate gyrus backprojection also provide a recurrent circuitry. Indeed, a study using computational modelling found that this backprojection has a crucial role in the ability of the computational model to perform simulated pattern separation and completion⁴⁹.

Novelty. It has been suggested that the dentate gyrus contributes to the ability to define what is familiar and what is novel in the environment^{88,89}. Mossy cells may contribute to this function by exciting granule cells when a novel sensory input is processed in the lateral entorhinal cortex and is sent to the dentate gyrus by the lateral perforant path. Such a pathway is suggested by recordings from a subset of mossy cells that have a low action potential threshold in response to a perforant path stimulus27. These mossy cells had dendrites in the molecular layer, where they could receive direct perforant path inputs²⁷. Whether a direct entorhinal cortex-mossy cell pathway exists has not been proven, but electrophysiological data suggest that perforant path axons make monosynaptic inputs onto mossy cells²⁷. Mossy cell dendrites in the molecular layer have been shown in diverse species²⁸, and in primates there are also dendrites of CA3c pyramidal cells in the molecular layer⁹⁰. Thus, mossy cells might act as sentinels that signal to granule cells that a change has occurred. This could ensure that the input from the entorhinal cortex always depolarizes some granule cells, even when it is too weak to reach the threshold for a granule cell action potential by itself.

Afterdepolarization A depolarization occurring after an event, typically after an action potential.

Another possible path from the entorhinal cortex to mossy cells involves the semilunar granule cells in the IML, which receive robust inputs from the entorhinal cortex and activate the mossy cells in a robust manner^{9,10}. Thus, an entorhinal cortex–semilunar granule cell–mossy cell pathway could explain the low action potential threshold of some mossy cells (FIG. 4d).

It is notable that the hilus receives afferents from the brainstem, such as inputs from the locus coeruleus^{76,91,92}. Many of these brainstem inputs are part of the reticular activating system and, therefore, would act in concert with entorhinal inputs to 'inform' the dentate gyrus about a changing environment. Mossy cells also receive cholinergic inputs⁹³; this is important because the activation of muscarinic cholinergic receptors depolarizes the mossy cells and elicits an afterdepolarization that does not occur in other hilar cell types⁹⁴. For these neuromodulatory inputs, mossy cells would potentially be activated even when other dentate gyrus neurons are not. Thus, the extrinsic afferents that preferentially activate the mossy cells may provide an alternative mechanism for a sentinel function.

In vivo support for the idea that mossy cells act as a type of sentinel comes from immunohistochemical studies of *Fos*, a marker of neuronal activity⁹⁵. When rats were removed from their home cage, brought to a laboratory and rapidly perfused, a subset of mossy cells in the ventral hippocampus expressed FOS protein, as did a subset of granule cells in the dorsal dentate gyrus (the target of ventral mossy cell projections)⁹⁵. Presumably, the expression of FOS in mossy cells occurred in response to small changes in the environment when the animals were in the home cage, or to rapid changes as the animals experienced novelty of the laboratory where they were brought to be perfused⁹⁵.

Anxiety. Behavioural studies using Crlr-Cre mice demonstrated that the diphtheria toxin-mediated deletion of mossy cells led to behaviours that suggested increased anxiety compared with controls¹. This might be expected, given that there are more mossy cells in the ventral dentate gyrus than in the dorsal dentate gyrus⁹⁶⁻⁹⁸ and given that lesions of the ventral dentate gyrus lead to anxiety99. One mechanism by which mossy cells might contribute to innate anxiety is related to the idea that mossy cells can signal changes in the environment (see above). Without such 'cues', there may be less certainty in the environment, such as certainty of safety from predators, leading to anxiety. Mossy cells may also play a part in other types of anxiety (as well as in fear and stress); indeed, transgenic mice in which mossy cells were ablated spent less time in the open arm of the elevated plus maze than control mice¹, a task that does not necessarily reflect innate anxiety.

Mossy cell vulnerability

A distinguishing feature of mossy cells is their vulnerability to insults or injury. Pathological studies show that the hilus is a common site of neuronal loss in temporal lobe epilepsy (TLE)¹⁰⁰, a subtype of epilepsy in which seizures involve the temporal lobe of the cortex. Although some mossy cells survive in TLE⁹⁷, many are vulnerable¹⁰¹. As animal models of TLE were developed, detailed comparisons of the cell types in the hilus showed that two hilar cell types were especially vulnerable: the mossy cells and the

Box 1 | Potential mechanisms of mossy cell vulnerability

Mossy cells may be vulnerable to insults or injury as a result of presynaptic mechanisms. For example, as shown in part **a** of the figure, the giant boutons of granule cells can release large concentrations of glutamate onto mossy cell thorny excrescences, resulting in excitotoxicity^{7,75,106,107,109}. Peptides such as brain-derived neurotrophic factor that are located in dense core vesicles within the giant boutons can facilitate the release of glutamate, exacerbating this excitotoxicity¹²⁵.

The vulnerability of mossy cells has also been suggested to arise owing to their weak expression of the calcium-binding proteins parvalbumin and calbindin, which results in poor buffering of intracellular calcium when it enters the cell²⁰. However, neurons outside the hippocampus that lack calcium-binding proteins are not necessarily vulnerable¹²⁶. Moreover, in a mouse model of epilepsy, mice that lack calcium-binding proteins have a phenotype similar to that of wild-type mice¹²⁷.

Mossy cells also express striatum-enriched

protein-tyrosine phosphatase (STEP), the expression of which has been suggested to explain the vulnerability of HIPP cells (hilar cells that project to the terminal zone of the perforant path)¹²⁸. Mossy cells express lower levels of STEP than HIPP cells (based on immunohistochemistry¹²⁸), but even a low expression level of STEP could contribute to their vulnerability given its importance for the vulnerability of HIPP cells¹²⁸. HIPP and other hilar cells, including mossy cells, also have low levels of the δ -subunit of type A GABA receptors, which normally contributes to GABAergic inhibition, and therefore low levels of this subunit could make a cell vulnerable to excitotoxicity¹²⁹.

It has been suggested that the vulnerability of mossy cells may be due to low levels of autophagy¹³⁰, the process involved in waste removal from neurons. In the absence of strong autophagy, it is possible that mossy cells may not be able to keep up with metabolic demands.

Finally, it has been suggested that the vulnerability of mossy cells may be due to their physiological properties. In the example shown in part **b** of the figure, a typical action potential (AP) in a granule cell that is evoked by an afferent input (arrow) gives rise to only one AP. This limited firing, attributed to the high threshold and strong spike frequency adaptation (SFA) of granule cells¹³¹, limits excitotoxicity. As shown, the AP generated in a basket cell is faster and is followed by strong afterhyperpolarization (AHP)^{13,16,17}, which also limits firing and excitotoxicity. However, an afferent input to a mossy cell leads more readily to APs, and the APs are of longer duration with relatively weak repolarization³⁵. Thus, excitotoxicity is more likely to occur in mossy cells. MF, mossy fibre.



somatostatin-expressing hilar interneurons (HIPP cells); in comparison, granule cells were more resistant^{7,19,21,101}. In rats, mossy cells die after many types of insults and injuries that are risk factors for TLE, including forebrain ischaemia^{7,102}, status epilepticus^{7,96-98} (with exceptions¹⁰³) and traumatic brain injury^{104,105}. This led to the hypothesis that a loss of hilar cells — and possibly a loss of mossy cells specifically — causes TLE¹⁰⁶. Some of the potential causes of mossy cell vulnerability^{75,107-109} are shown in BOX 1.

The hypothesis that a loss of mossy cells causes TLE remains a subject of debate. One argument for this hypothesis notes that mossy cells can inhibit granule cells, which may normally prevent seizure activity from passing from the entorhinal cortex to the hippocampus¹¹⁰. Entry of seizure activity into the hippocampus could be important to sustain and amplify seizures. It has also been proposed that mossy cell-mediated excitation of granule cells is increased after insults that lead to TLE, causing hyperexcitability^{1,104–106,111,112}. However, hyperexcitability

could also be caused by a loss of HIPP cells⁹⁷ or calretininexpressing interneurons⁷⁰. Diverse changes in other aspects of the inhibitory control of granule cells¹¹³, including changes in the subunit composition of type A GABA receptors¹¹⁴⁻¹¹⁶, have led to the current consensus that a loss of mossy cells may contribute to TLE but is unlikely to be fully responsible for it.

One limitation of animal models of TLE is that many changes occur in response to the induction of epilepsy, not only a loss of mossy cells. Because of the relative specificity of the deletion of mossy cells in *Crlr*–Cre mice treated with the diptheria toxin, it was important to note that the mice showed no spontaneous seizures¹, supporting the idea that a loss of mossy cells is insufficient to cause TLE. However, the mice did exhibit transient disinhibition in response to perforant path stimulation, showing that the loss of mossy cells did have a marked effect in these mice. In addition, the mouse strain used to generate the *Crlr*–Cre mice was seizure resistant, as noted by the authors⁶¹.

Spike frequency adaptation (SFA). A reduction in frequency of action potential discharge during a constant depolarizing input. Most neurons have a high firing frequency after the beginning of a strong depolarization, and the firing frequency decays if the depolarization continues.

Contextual discrimination A behavioural task that tests the ability to distinguish two environments or contexts.

How a loss of mossy cells would contribute to TLE is unknown. One possibility is that the loss of mossy cells could make seizures more severe. In support of this idea, mice lacking mossy cells had more severe seizures than controls¹. Interestingly, the deletion of adult-born granule cells also increases the severity of seizures¹¹⁷. Thus, mossy cells could have an important role in TLE through their activation of immature adult-born granule cells, which innervate GABAergic neurons to a greater extent than excitatory neurons until their giant boutons develop¹¹⁸.

Mossy cells may also influence other diseases. For example, mossy cells and basket cells express p11 and helicase-like transcription factor (SMARCA3), proteins that are potentially important in major depressive disorder¹¹⁹. Epigenetic changes mediated by p11 through its actions on SMARCA3 (REF. 119) could change the physiological properties of mossy cells and basket cells, making their contribution to dentate gyrus function more robust or weaker. Mossy cells also express dysbindin 1C120 and dopamine D2 receptors^{59,60}, both of which are implicated in schizophrenia. The fact that a loss of mossy cells causes impaired contextual discrimination1, and that mice without mossy cells show behaviours that are consistent with increased anxiety (described above), suggests that mossy cells may have roles in psychiatric disorders. Precisely what these roles are remains to be determined.

Conclusions and perspectives

Experiments that have preferentially deleted or activated mossy cells have supported the long-standing hypothesis that the inhibition of granule cells is the major functional role of mossy cells. However, experiments in awake behaving animals are still lacking. Without such physiological data one wonders whether the conditions for detection of mossy cell-mediated excitation of granule cells have simply not been met. There is a gap between studies of dentate gyrus circuitry and studies of dentate gyrusdependent behaviour, and a major goal for the future would be to fill this void. Current approaches to address this issue are labour intensive and expensive, but rewarding. For example, patch clamp recordings in the dentate gyrus of awake, head-fixed mice have been made121; if mossy cells can be patched, a great deal of information could be gleaned. A more feasible alternative may be to use juxtacellular recording122, which allows many questions to be addressed, particularly when it is combined with electrodes to monitor other hippocampal subfields or oscillatory behaviour. Microendoscopy to image the hilus in vivo in awake behaving mice is becoming feasible¹²³. Using dyes such as GCaMP6 (REF. 124), expressed in cells under the control of Cre, would make selective imaging of mossy cells in vivo possible. However, these techniques require significant trauma to overlying tissues, to which mossy cells are particularly sensitive. While investigators attempt to circumvent this and other difficulties, other fundamental questions also need to be addressed. For example, quantitative data about mossy cell inputs and their outputs are surprisingly incomplete for rodents and especially for primates, for both sexes and for diverse ages. Such data would help to refine hypotheses, fuel the development of approaches such as computational modelling and ultimately make mossy cells much less enigmatic than they are now.

 Jinde, S. *et al.* Hilar mossy cell degeneration causes transient dentate granule cell hyperexcitability and impaired pattern separation. *Neuron* 76, 1189–1200 (2012).

This study examined the consequence of ablating mossy cells with the most selective approach to date. Although some CA3 pyramidal cells were also affected, the results showed that mossy cells have an important role in the dentate gyrus network.

- Hsu, T. T., Lee, C. T., Tai, M. H. & Lien, C. C. Differential recruitment of dentate gyrus interneuron types by commissural versus perforant pathways. *Cereb. Cortex* 26, 2715–2727 (2016).
 This study used optogenetics in hippocampal slices to show that mossy cells could excite or inhibit granule cells; the authors found that the major effect was inhibition.
- Amaral, D.G., Scharfman, H. E. & Lavenex, P. The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Prog. Brain Res.* 163, 3–22 (2007).
- Steward, O. & Scoville, S. A. Cells of origin of entorhinal cortical afferents to the hippocampus and fascia dentata of the rat. *J. Comp. Neurol.* **169**, 347–370 (1976).
- Witter, M. P. The perforant path: projections from the entorhinal cortex to the dentate gyrus. *Prog. Brain Res.* 163, 43–61 (2007).
- Ribak, C. E., Seress, L. & Amaral, D. G. The development, ultrastructure and synaptic connections of the mossy cells of the dentate gyrus. *J. Neurocytol.* 14, 835–857 (1985).
- Scharfman, H. E. & Myers, C. E. Hilar mossy cells of the dentate gyrus: a historical perspective. *Front. Neural Circuits* 6, 106 (2012).
- Buckmaster, P. S., Wenzel, H. J., Kunkel, D. D. & Schwartzkroin, P. A. Axon arbors and synaptic connections of hippocampal mossy cells in the rat *in vivo. J. Comp. Neurol.* 366, 271–292 (1996).

This paper is an excellent quantitative study of the mossy cell projection *in vivo*.

- Larimer, P. & Strowbridge, B. W. Representing information in cell assemblies: persistent activity mediated by semilunar granule cells. *Nat. Neurosci.* 13, 213–222 (2010).
- Williams, P. A., Larimer, P., Gao, Y. & Strowbridge, B. W. Semilunar granule cells: glutamatergic neurons in the rat dentate gyrus with axon collaterals in the inner molecular layer. *J. Neurosci.* 27, 13756–13761 (2007). This study was the first to characterize semilunar granule cells with electrophysiology, and it showed their potential importance in the network of the dentate gyrus.
- Scharfman, H. E., Goodman, J. & McCloskey, D. Ectopic granule cells of the rat dentate gyrus. *Dev. Neurosci.* 29, 14–27 (2007).
- Kempermann, G., Song, H. & Gage, F. H. Neurogenesis in the adult hippocampus. *Cold Spring Harb. Perspect. Biol.* 7, a018812 (2015).
- 13. Freund, T. F., Buzsaki, G. Interneurons of the hippocampus. *Hippocampus* **6**, 347–470 (1996).
- Halasy, K. & Somogyi, P. Subdivisions in the multiple GABAergic innervation of granule cells in the dentate gyrus of the rat hippocampus. *Eur. J. Neurosci.* 5, 411–429 (1993).
- 15. Han, Z. S., Buhl, E. H., Lorinczi, Z. & Somogyi, P. A high degree of spatial selectivity in the axonal and dendritic domains of physiologically identified localcircuit neurons in the dentate gyrus of the rat hippocampus. *Eur. J. Neurosci.* 5, 395–410 (1993). References 14 and 15 proposed an organization of the dentate gyrus interneurons according to the location of the cell body and terminal field of the axon.
- Hosp, J. A. *et al.* Morpho-physiological criteria divide dentate gyrus interneurons into classes. *Hippocampus* 24, 189–203 (2014).

This paper suggested an alternative nomenclature to the organization of interneurons proposed in references 14 and 15 that reconciled discrepancies in the literature.

- Scharfman, H. E. Electrophysiological diversity of pyramidal-shaped neurons at the granule cell layer/ hilus border of the rat dentate gyrus recorded *in vitro*. *Hippocampus* 5, 287–305 (1995).
- Seay-Lowe, S. L. & Claiborne, B. J. Morphology of intracellularly labeled interneurons in the dentate gyrus of the immature rat. *J. Comp. Neurol.* **324**, 23–36 (1992).
- Sloviter, R. S. Decreased hippocampal inhibition and a selective loss of interneurons in experimental epilepsy. *Science* 235, 73–76 (1987).
 This paper showed that mossy cells were vulnerable to seizures but that many GABAergic neurons were not, which was surprising because a loss of GABAergic neurons was considered to be a cause of epilepsy.
- Sloviter, R. S. Calcium-binding protein (calbindin-D_{28k}) and parvalbumin immunocytochemistry: localization in the rat hippocampus with specific reference to the selective vulnerability of hippocampal neurons to seizure activity. *J. Comp. Neurol.* **280**, 183–196 (1989).
- Schwarzer, C., Williamson, J. M., Lothman, E. W., Vezzani, A. & Sperk, G. Somatostatin, neuropeptide Y, neurokinin B and cholecystokinin immunoreactivity in two chronic models of temporal lobe epilepsy. *Neuroscience* 69, 831–845 (1995).
- Houser, C. R. Interneurons of the dentate gyrus: an overview of cell types, terminal fields and neurochemical identity. *Prog. Brain Res.* 163, 217–232 (2007).
- Goodman, J. H. & Sloviter, R. S. Evidence for commissurally projecting parvalbumin-immunoreactive basket cells in the dentate gyrus of the rat. *Hippocampus* 2, 13–21 (1992).

- 24. Deller, T., Nitsch, R. & Frotscher, M. Phaseolus vulgaris-leucoagglutinin tracing of commissural fibers to the rat dentate gyrus: evidence for a previously unknown commissural projection to the outer molecular layer. J. Comp. Neurol. 352, 55-68 (1995).
- Amaral, D. G. A golgi study of cell types in the hilar 25. region of the hippocampus in the rat. J. Comp. Neurol. 15, 851-914 (1978).

This paper provided the first detailed description of hilar cells and mossy cells. Frotscher, M., Seress, L., Schwerdtfeger, W. K. &

- 26. Buhl, E. The mossy cells of the fascia dentata: a comparative study of their fine structure and synaptic connections in rodents and primates. J Comp. Neurol. **312**, 145–163 (1991).
- Scharfman, H. E. Dentate hilar cells with dendrites in the molecular layer have lower thresholds for synaptic activation by perforant path than granule cells. *J. Neurosci.* **11**, 1660–1673 (1991). Blackstad, J. B. *et al.* Observations on hippocampal
- 28 mossy cells in mink (neovison vison) with special reference to dendrites ascending to the granular and molecular layers. Hippocampus 26, 229-245 (2016).
- 29 Chicurel, M. E. & Harris, K. M. Three-dimensional analysis of the structure and composition of CA3 branched dendritic spines and their synaptic relationships with mossy fiber boutons in the rat hippocampus. J. Comp. Neurol. 325, 169–182 (1992).
- 30 Blackstad, T. W. & Kjaerheim, A. Special axo-dendritic synapses in the hippocampal cortex: electron and light microscopic studies on the layer of mossy fibers. J. Comp. Neurol. 117, 133–159 (1961).
- Henze, D. A., Urban, N. N. & Barrionuevo, G. 31. The multifarious hippocampal mossy fiber pathway: a review. *Neuroscience* **98**, 407–427 (2000). Laatsch, R. H. & Cowan, W. M. Electron microscopic
- 32 studies of the dentate gyrus of the rat. I. Normal structure with special reference to synaptic organization. J. Comp. Neurol. 128, 359–395 (1966).
- Amaral, D. G. & Dent, J. A. Development of the mossy fibers of the dentate gyrus: I. A light and 33 electron microscopic study of the mossy fibers and their expansions. J. Comp. Neurol. 195, 51-86 (1981).
- 34. Acsady, L., Kamondi, A., Sik, A., Freund, T. & Buzsaki, G. GABAergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. J. Neurosci. 18, 3386–3403 (1998). Using quantitative anatomical methods, this paper identified that mossy fibre synapses on GABAergic neurons outnumber those on CA3 pyramidal cells, suggesting that the pathway has strong inhibitory effects.
- Scharfman, H. E. & Schwartzkroin, P. A 35. Electrophysiology of morphologically identified mossy cells of the dentate hilus recorded in guinea pig hippocampal slices. J. Neurosci. 8, 3812-3821 (1988)
- Scharfman, H. E. Characteristics of spontaneous and 36 evoked EPSPs recorded from dentate spiny hilar cells in rat hippocampal slices. J. Neurophysiol. 70, 742-757 (1993)
- Seress, L., Abraham, H., Doczi, T., Lazar, G. & Kozicz, T. Cocaine- and amphetamine-regulated 37. transcript peptide (CART) is a selective marker of rat granule cells and of human mossy cells in the hippocampal dentate gyrus. Neuroscience 125, 13-24 (2004).

This paper showed that there are important exceptions to the idea that mossy cells are always lost in TLE.

- 38 Blasco-Ibanez, J. M. & Freund, T. F. Distribution, ultrastructure, and connectivity of calretinin-immunoreactive mossy cells of the mouse dentate gyrus. Hippocampus 7, 307–320 (1997).
- Fujise, N., Liu, Y., Hori, N. & Kosaka, T. Distribution of 39 calretinin immunoreactivity in the mouse dentate gyrus: II. Mossy cells, with special reference to their dorsoventral difference in calretinin immunoreactivity. Neuroscience 82, 181–200 (1998).
- 40. Zimmer, J. Ipsilateral afferents to the commissural zone of the fascia dentata, demonstrated in decommissurated rats by silver impregnation
- J. Comp. Neurol. 142, 393–416 (1971). Berger, T. W., Semple-Rowland, S. & Bassett, J. L. 41 Hippocampal polymorph neurons are the cells of origin for ipsilateral association and commissural afferents to the dentate gyrus. Brain Res. 224, 329-336 (1981).

- 42. Buckmaster, P. S., Strowbridge, B. W., Kunkel, D. D., Schmiege, D. L. & Schwartzkroin, P. A. Mossy cell axonal projections to the dentate gyrus molecular laver in the rat hippocampal slice. Hippocampus 2. 349-362 (1992).
- Moser, M. B. & Moser, E. I. Functional differentiation 43. in the hippocampus. Hippocampus 8, 608-619 (1998).
- 44 Sahav A Drew M R & Hen R Dentate gyrus neurogenesis and depression. Prog. Brain Res. 163, 697-722 (2007).
- Phillips, R. G. & LeDoux, J. E. Differential contribution 45. of amygdala and hippocampus to cued and contextual fear conditioning. Behav. Neurosci. 106, 274-285 (1992).
- 46. Kesner, R. P. A behavioral analysis of dentate gyrus function. Prog. Brain Res. 163, 567-576 (2007).
- 47. Rolls, E. T. Pattern separation, completion, and categorisation in the hippocampus and neocortex. Neurobiol. Learn. Mem. **129**, 4–28 (2016). Yassa, M. A. & Stark, C. E. Pattern separation in the
- 48. hippocampus. Trends Neurosci. 34, 515-525 (2011)
- 49. Myers, C. E. & Scharfman, H. E. Pattern separation in the dentate gyrus: a role for the CA3 backprojection. *Hippocampus* **21**, 1190–1215 (2011). Myers, C. E. & Scharfman, H. E. A role for hilar cells in
- 50 pattern separation in the dentate gyrus: a computational approach. Hippocampus 19, 321-337 (2009)
- Buzsaki, G. & Eidelberg, E. Commissural projection to 51. the dentate gyrus of the rat: evidence for feed-forward inhibition. *Brain Res.* **230**, 346–350 (1981).
- Douglas, R. M., McNaughton, B. L. & Goddard, G. V. 52. Commissural inhibition and facilitation of granule cell discharge in fascia dentata. J. Comp. Neurol. 219, 285-294 (1983).
- *Hippocampus* **4**, 65–69 (1994). This paper provided the first evidence that mossy cells were glutamatergic.
- Scharfman, H. E. Electrophysiological evidence that 54 dentate hilar mossy cells are excitatory and innervate both granule cells and interneurons. J. Neurophysiol. **74**, 179–194 (1995).
- This paper showed that mossy cells directly excite granule cells and dentate gyrus interneurons 55
- Larimer, P. & Strowbridge, B. W. Nonrandom local circuits in the dentate gyrus. *J. Neurosci.* 28, 12212–12223 (2008).
- 56 Ratzliff, A. d. H., Howard, A. L., Santhakumar, V., Osapay, I. & Soltesz, I. Rapid deletion of mossy cells does not result in a hyperexcitable dentate gyrus: implications for epileptogenesis. J. Neurosci. 24, 2259-2269 (2004).

This paper used a method to delete mossy cells in a hippocampal slice and did not find any evidence for hyperexcitability, thus arguing against the idea that mossy cells normally activate interneurons.

- Jackson, M. B. & Scharfman, H. E. Positive feedback 57. from hilar mossy cells to granule cells in the dentate gyrus revealed by voltage-sensitive dye and microelectrode recording. J. Neurophysiol. 76, 601-616 (1996).
- 58. Wright, B. J. & Jackson, M. B. Long-term potentiation in hilar circuitry modulates gating by the dentate gyrus. J. Neurosci. 34, 9743-9753 (2014).
- Gangarossa, G. et al. Characterization of dopamine D1 59. and D2 receptor-expressing neurons in the mouse hippocampus. *Hippocampus* **22**, 2199–2207 (2012). Puighermanal, E. *et al. drd2-cre:ribotag* mouse line
- 60. unravels the possible diversity of dopamine D2 receptor-expressing cells of the dorsal mouse hippocampus. Hippocampus 25, 858-875 (2015).
- 61. Jinde, S., Zsiros, V. & Nakazawa, K. Hilar mossy cell circuitry controlling dentate granule cell excitability. Front. Neural Circuits 7, 14 (2013).
- Chancey, J. H., Poulsen, D. J., Wadiche, J. I. & 62. Overstreet-Wadiche, L. Hilar mossy cells provide the first glutamatergic synapses to adult-born dentate granule cells. J. Neurosci. 34 2349-2354 (2014).
- Lysetskiy, M., Foldy, C. & Soltesz, I. Long- and shortterm plasticity at mossy fiber synapses on mossy cells in the rat dentate gyrus. Hippocampus 15, 691-696 (2005)
- Hetherington, P. A., Austin, K. B. & Shapiro, M. L. 64 Ipsilateral associational pathway in the dentate gyrus: an excitatory feedback system that supports N-methyl-p-aspartate-dependent long-term potentiation. Hippocampus 4, 422-438 (1994).

- 65. Kleschevnikov, A. M. & Routtenberg, A. Long-term potentiation recruits a trisynaptic excitatory associative network within the mouse dentate evrus. *Eur. J. Neurosci.* **17**, 2690–2702 (2003).
- 66 Alvarez-Salvado, E., Pallares, V., Moreno, A. & Canals, S. Functional MRI of long-term potentiation: imaging network plasticity. *Phil. Trans. R. Soc. B* **369**, 20130152 (2014).
- 67. Chiu, C. Q. & Castillo, P. E. Input-specific plasticity at excitatory synapses mediated by endocannabinoids in the dentate gyrus. *Neuropharmacology* **54**, 68–78 (2008)
- Hofmann, M. E., Nahir, B. & Frazier, C. J. 68. Endocannabinoid-mediated depolarization-induced suppression of inhibition in hilar mossy cells of the rat dentate gyrus. J. Neurophysiol. **96**, 2501–2512 (2006).
- Schurmans, S. et al. Impaired long-term potentiation 69 induction in dentate gyrus of calretinin-deficient mice. *Proc. Natl Acad. Sci. USA* **94**, 10415–10420 (1997). Töth, K. & Maglóczky, Z. The vulnerability of calretinin-
- 70. containing hippocampal interneurons to temporal lobe epilepsy. Front. Neuroanat. 8, 100 (2014).
- Namgung, U., Matsuyama, S. & Routtenberg, A Long-term potentiation activates the GAP-43 promoter: selective participation of hippocampal mossy cells. *Proc. Natl Acad. Sci. USA* **94**, 11675–11680 (1997). Buzsaki, G. & Moser, E. I. Memory, navigation and
- 72. theta rhythm in the hippocampal-entorhinal system. Nat. Neurosci. 16, 130-138 (2013).
- Soltesz, I., Bourassa, J. & Deschenes, M. The behavior of mossy cells of the rat dentate gyrus during theta 73 oscillations in vivo. Neuroscience 57, 555-564 (1993).
- 74 Henze, D. A. & Buzsáki, G. Hilar mossy cells: functional identification and activity in vivo. *Prog. Brain Res.* **163**, 199–216 (2007).
- Buckmaster, P. S. & Schwartzkroin, P. A. Hippocampal 75. mossy cell function: a speculative view. Hippocampus 4, 393-402 (1994).
- Brown, R. A., Walling, S. G., Milway, J. S. & Harley, C. W. Locus ceruleus activation suppresses feedforward 76 interneurons and reduces β - γ electroencephalogram frequencies while it enhances 0 frequencies in rat dentate gyrus. *J. Neurosci.* **25**, 1985–1991 (2005). Sahay, A., Wilson, D. A. & Hen, R. Pattern
- 77 separation: a common function for new neurons in hippocampus and olfactory bulb. Neuron 70, 582-588 (2011).
- Aimone, J. B., Deng, W. & Gage, F. H. Resolving new 78 memories: a critical look at the dentate gyrus, adult neurogenesis, and pattern separation. Neuron 70, 589-596 (2011)
- Alme, C. B. et al. Hippocampal granule cells opt for early 79. retirement. *Hippocampus* **20**, 1109–1123 (2010).
- 80 Scharfman, H. E. The CA 3 "backprojection" to the dentate gyrus. *Prog. Brain Res.* **163**, 627–637 (2007). Li, X., Somogyi, P., Ylinen, A. & Buzsáki, G. The 81
- hippocampal CA3 network: an in vivo intracellular labeling study. J. Comp. Neurol. 339, 181-208 (1994).
- 82 Ishizuka, N., Weber, J. & Amaral, D. G. Organization of intrahippocampal projections originating from CA3 pyramidal cells in the rat. J. Comp. Neurol. 295 580-623 (1990).

References 81 and 82 showed that there was a pattern of innervation 'back' to the dentate gyrus, opposite to the direction of the trisynaptic circuitry.

- Scharfman, H. E. EPSPs of dentate gyrus granule cells 83 during epileptiform bursts of dentate hilar "mossy" cells and area CA3 pyramidal cells in disinhibited rat hippocampal slices. J. Neurosci. 14, 6041–6057 (1994)
- 84 Scharfman, H. E. Evidence from simultaneous intracellular recordings in rat hippocampal slices that area CA3 pyramidal cells innervate dentate hilar mossy cells. J. Neurophysiol. 72, 2167-2180 (1994). References 83 and 84 showed the physiological effects that supported the idea that pyramidal cells innervate hilar neurons such as mossy cells.
- Kneisler, T. B. & Dingledine, R. Synaptic input from 85 CA3 pyramidal cells to dentate basket cells in rat hippocampus. *J. Physiol.* **487**, 125–146 (1995). Lisman, J. E., Talamini, L. M. & Raffone, A. Recall of
- 86 memory sequences by interaction of the dentate and CA3: a revised model of the phase precession. Neural Netw. 18, 1191-1201 (2005).
- Penttonen, M., Kamondi, A., Sik, A., Acsady, L. & 87. Buzsaki, G. Feed-forward and feed-back activation of the dentate gyrus in vivo during dentate spikes and
- sharp wave bursts. *Hippocampus* **7**, 437–450 (1997). Hunsaker, M. R., Rosenberg, J. S. & Kesner, R. P. 88. The role of the dentate gyrus, CA3a,b, and CA3c for detecting spatial and environmental novelty. Hippocampus 18, 1064-1073 (2008).

Soriano, E. & Frotscher, M. Mossy cells of the rat fascia dentata are glutamate-immunoreactive. 53

- Aggleton, J. P., Brown, M. W. & Albasser, M. M. Contrasting brain activity patterns for item recognition memory and associative recognition memory: insights from immediate-early gene functional imaging. *Neuropsychologia* 50, 3141–3155 (2013).
- Buckmaster, P. S. & Amaral, D. G. Intracellular recording and labeling of mossy cells and proximal CA3 pyramidal cells in macaque monkeys. *J. Comp. Neurol.* **430**, 264–281 (2001).
 Amaral, D. G. & Campbell, M. J. Transmitter systems
- Amaral, D. G. & Campbell, M. J. Transmitter systems in the primate dentate gyrus. *Hum. Neurobiol.* 5, 169–180 (1986).
- Swanson, L. W., Kohler, C. & Bjorklund, A. in Handbook of Chemical Neuroanatomy Vol. 5 (eds Hokfelt, T., Bjorklund, A. & Swanson, L. W.) 125–277 (Elsevier, 1987).
- Deller, T., Katona, I., Cozzari, C., Frotscher, M. & Freund, T. F. Cholinergic innervation of mossy cells in the rat fascia dentata. *Hippocampus* 9, 314–320 (1999).
- Hofmann, M. E. & Frazier, C. J. Muscarinic receptor activation modulates the excitability of hilar mossy cells through the induction of an afterdepolarization. *Brain Res.* 1318, 42–51 (2010).
- Duffy, A. M., Schaner, M. J., Chin, J. & Scharfman, H. E. Expression of c-fos in hilar mossy cells of the dentate gyrus *in vivo*. *Hippocampus* 23, 649–655 (2013).
- Jiao, Y. & Nadler, J. V. Stereological analysis of GluR2-immunoreactive hilar neurons in the pilocarpine model of temporal lobe epilepsy: correlation of cell loss with mossy fiber sprouting. *Exp. Neurol.* 205, 569–582 (2007).
- Buckmaster, P. S. & Jongen-Relo, A. L. Highly specific neuron loss preserves lateral inhibitory circuits in the dentate gyrus of kainate-induced epileptic rats. *J. Neurosci.* **19**, 9519–9529 (1999).
- Volz, F. *et al.* Stereologic estimation of hippocampal GluR2/3- and calretinin-immunoreactive hilar neurons (presumptive mossy cells) in two mouse models of temporal lobe epilepsy. *Epilepsia* 52, 1579–1589 (2011).
- Bannerman, D. M. *et al.* Ventral hippocampal lesions affect anxiety but not spatial learning. *Behav. Brain Res.* 139, 197–213 (2003).
- Margerison, J. & Corsellis, J. Epilepsy and the temporal lobes: a clinical, electrographic and neuropathological study of the brain in epilepsy with particular reference to the temporal lobes. *Brain* 89, 499–530 (1966).
- 101. de Lanerolle, N. C., Lee, T. S. & Spencer, D. D. in Jasper's Basic Mechanisms of the Epilepsies 4th edn (eds Noebels, J. L., Avoli, M., Rogawski, M. A., Olsen, R. W. & Delgado-Escueta, A. V.) 387–404 (Oxford Univ. Press, 2012)
- 102. Crain, B. J., Westerkam, W. D., Harrison, A. H. & Nadler, J. V. Selective neuronal death after transient forebrain ischemia in the mongolian gerbil: a silver impreenation study. *Neuroscience* 27, 387–402 (1988)
- impregnation study. Neuroscience 27, 387–402 (1988).
 103. Kotti, T., Tapiola, T., Riekkinen, P. J. Sr. & Miettinen, R. The calretinin-containing mossy cells survive excitotoxic insult in the gerbil dentate gyrus. Comparison of excitotoxicity-induced neuropathological changes in the gerbil and rat. *Eur. J. Neurosci.* 8, 2371–2378 (1996).
 104. Lowenstein, D. H., Thomas, M. J., Smith, D. H. &
- 104. Lowenstein, D. H., Thomas, M. J., Smith, D. H. & McIntosh, T. K. Selective vulnerability of dentate hilar neurons following traumatic brain injury: a potential mechanistic link between head trauma and disorders of the hippocampus. *J. Neurosci.* **12**, 4846–4853 (1992).
- 105. Santhakumar, V. *et al.* Granule cell hyperexcitability in the early post-traumatic rat dentate gyrus: the 'irritable mossy cell' hypothesis. *J. Physiol.* 524, 117–134 (2000).
 This paper showed that after head trauma there is hyperexcitability in the dentate gyrus and that

hyperexcitability in the dentate gyrus and that mossy cells may contribute to the hyperexcitability because they become activated or 'irritable' and therefore excite granule cells more than normal.

- 106. Sloviter, R. S. The functional organization of the hippocampal dentate gyrus and its relevance to the pathogenesis of temporal lobe epilepsy. *Ann. Neurol.* **35**, 640–654 (1994).
- 107. Scharfman, H. E. The role of nonprincipal cells in dentate gyrus excitability and its relevance to animal models of epilepsy and temporal lobe epilepsy. *Adv. Neurol.* **79**, 805–820 (1999).
- 108. Tsankova, N. M., Kumar, A. & Nestler, E. J. Histone modifications at gene promoter regions in rat hippocampus after acute and chronic electroconvulsive seizures. J. Neurosci. 24, 5603–5610 (2004).

- 109. Schwartzkroin, P. A. *et al.* Possible mechanisms of seizure-related cell damage in the dentate hilus. *Epilepsy Res. Suppl.* **12**, 317–324 (1996).
- Hsu, D. The dentate gyrus as a filter or gate: a look back and a look ahead. *Prog. Brain Res.* 163, 601–613 (2007).
- Ratzliff, A., Santhakumar, V., Howard, A. & Soltesz, I. Mossy cells in epilepsy: rigor mortis or vigor mortis? *Trends Neurosci.* 25, 140–144 (2002).
- 112. Sloviter, R. S. Permanently altered hippocampal structure, excitability, and inhibition after experimental status epilepticus in the rat: the "dormant basket cell" hypothesis and its possible relevance to temporal lobe epilepsy. *Hippocampus* 1, 41–66 (1991).
- Zhang, W. & Buckmaster, P. S. Dysfunction of the dentate basket cell circuit in a rat model of temporal lobe epilepsy. J. Neurosci. 29, 78/6–7856 (2009)
- 7846–7856 (2009).
 114. Brooks-Kayal, A. R., Shumate, M. D., Jin, H., Rikhter, T. Y. & Coulter, D. A. Selective changes in single cell GABA, receptor subunit expression and function in temporal lobe epilepsy. *Nat. Med.* 4, 1166–1172 (1998).
- Coulter, D. A. Epilepsy-associated plasticity in γ-aminobutyric acid receptor expression, function, and inhibitory synaptic properties. *Int. Rev. Neurobiol.* 45, 237–252 (2001).
- 116. Sperk, G., Furtinger, S., Schwarzer, C. & Pirker, S. GABA and its receptors in epilepsy. *Adv. Exp. Med. Biol.* 548, 92–103 (2004). References 113–116 describe complex changes to GABAergic inhibition in the dentate gyrus in models of TLE and that one mechanism, such as a loss of mossy cell input to interneurons, is unlikely to explain the pathophysiology of TLE.
- Ivengar, S. S. *et al.* Suppression of adult neurogenesis increases the acute effects of kainic acid. *Exp. Neurol.* 264, 135–149 (2015).
- Scharfman, H. E. & Bernstein, H. L. Potential implications of a monosynaptic pathway from mossy cells to adult-born granule cells of the dentate gyrus. *Front. Syst. Neurosci.* 9, 112 (2015).
- Oh, Y. S. *et al.* SMARCA3, a chromatin-remodeling factor, is required for p11-dependent antidepressant action. *Cell* **152**, 831–843 (2013).
- 120. Wang, H. *et al.* Dysbindin-1C is required for the survival of hilar mossy cells and the maturation of adult newborn neurons in dentate gyrus. *J. Biol. Chem.* 289, 29060–29072 (2014).
- Anderson, R. W. & Strowbridge, B. W. α-Band oscillations in intracellular membrane potentials of dentate gyrus neurons in awake rodents. *Learn. Mem.* 21, 656–661 (2014).
- Tang, Q., Brecht, M. & Burgalossi, A. Juxtacellular recording and morphological identification of single neurons in freely moving rats. *Nat. Protoc.* 9, 2369–2381 (2014).
 Barretto, R. P. & Schnitzer, M. J. *In vivo*
- 123. Barretto, R. P. & Schnitzer, M. J. In vivo microendoscopy of the hippocampus. Cold Spring Harb. Protoc. 2012, 1092–1099 (2012).
- 124. Chen, T. W. et al. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499, 295–300 (2013).
- 125. Scharfman, H. E. in Synaptic Plasticity and Transsynaptic Signalling (eds Stanton, P. K., Bramham, C. R. & Scharfman, H. E.) 201–220 (Springer, 2005).
- 126. Freund, T. F., Buzsaki, G., Leon, A., Baimbridge, K. G. & Somogyi, P. Relationship of neuronal vulnerability and calcium binding protein immunoreactivity in ischemia. *Exp. Brain Res.* 83, 55–66 (1990).
- 127. Bouilleret, V., Schwaller, B., Schurmans, S., Celio, M. R. & Fritschy, J. M. Neurodegenerative and morphogenic changes in a mouse model of temporal lobe epilepsy do not depend on the expression of the calcium-binding proteins parvalbumin, calbindin, or calretinin. *Neuroscience* **97**, 47–58 (2000).
- 128. Choi, Y. S. et al. Status epilepticus-induced somatostatinergic hilar interneuron degeneration is regulated by striatal enriched protein tyrosine phosphatase. J. Neurosci. 27, 2999–3009 (2007).
- 129. Tong, X. et al. Ectopic expression of α6 and δ GABA_A receptor subunits in hilar somatostatin neurons increases tonic inhibition and alters network activity in the dentate gyrus. J. Neurosci. **35**, 16142–16158 (2015).
- 130. Yuan, Y., Wang, H., Wei, Z. & Li, W. Impaired autophagy in hilar mossy cells of the dentate gyrus and its implication in schizophrenia. *J. Genet. Genom* 42, 1–8 (2015).

- Staley, K. J., Otis, T. S. & Mody, I. Membrane properties of dentate gyrus granule cells: comparison of sharp microelectrode and whole-cell recordings. *J. Neurophysiol.* 67, 1346–1358 (1992).
- Etter, G. & Krezel, W. Dopamine D2 receptor controls hilar mossy cells excitability. *Hippocampus* 24, 725–732 (2014).
- 133. Leranth, C. & Frotscher, M. Cholinergic innervation of hippocampal GAD- and somatostatin-immunoreactive commissural neurons. J. Comp. Neurol. 261, 33–47 (1987).
- Ribak, C. E. Local circuitry of GABAergic basket cells in the dentate gyrus. *Epilepsy Res. Suppl.* 7, 29–47 (1992).
- 135. Armstrong, C., Szabadics, J., Tamas, G. & Soltesz, I. Neurogliaform cells in the molecular layer of the dentate gyrus as feed-forward *p*-aminobutyric acidergic modulators of entorhinal-hippocampal interplay. *J. Comp. Neurol.* **519**, 1476–1491 (2011).
- Gulyas, A. I., Hajos, N. & Freund, T. F. Interneurons containing calretinin are specialized to control other interneurons in the rat hippocampus. *J. Neurosci.* 16, 3397–3411 (1996).
- Freund, T. F. GABAergic septal and serotonergic median raphe afferents preferentially innervate inhibitory interneurons in the hippocampus and dentate gyrus. *Epilepsy Res. Suppl.* **7**, 79–91 (1992).
 Leranth, C. & Hajszan, T. Extrinsic afferent systems to
- the dentate gyrus. *Prog. Brain Res.* **163**, 63–84 (2007).
- 139. Wyss, J. M., Swanson, L. W. & Cowan, W. M. Evidence for an input to the molecular layer and the stratum granulosum of the dentate gyrus from the supramammillary region of the hypothalamus. *Anat. Embryol. (Berl.)* **156**, 165–176 (1979).
- 140. Toni, N. et al. Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nat. Neurosci.* **11**, 901–907 (2008).
- 141. Brandt, M. D. et al. Transient calretinin expression defines early postmitotic step of neuronal differentiation in adult hippocampal neurogenesis of mice. Mol. Cell. Neurosci. 24, 603–613 (2003).
- mice. Mol. Cell. Neurosci. 24, 603–613 (2003).
 142. Leranth, C., Szeidemann, Z., Hsu, M. & Buzsáki, G. AMPA receptors in the rat and primate hippocampus: a possible absence of GluR2/3 subunits in most interneurons. Neuroscience 70, 631–652 (1996).
- 143. Liu, Y., Fujise, N. & Kosaka, T. Distribution of calretinin immunoreactivity in the mouse dentate gyrus. I. General description. *Exp. Brain Res.* **108**, 389–403 (1996).
- Freund, T. F., Hajos, N., Acsady, L., Gorcs, T. J. & Katona, I. Mossy cells of the rat dentate gyrus are immunoreactive for calcitonin gene-related peptide (CGRP). *Eur. J. Neurosci.* 9, 1815–1830 (1997).
 Patel, A. & Bulloch, K. Type II glucocorticoid receptor
- Patel, A. & Bulloch, K. Type II glucocorticoid receptor immunoreactivity in the mossy cells of the rat and the mouse hippocampus. *Hippocampus* 13, 59–66 (2003).
- 146. Lubke, J., Frotscher, M. & Spruston, N. Specialized electrophysiological properties of anatomically identified neurons in the hilar region of the rat fascia dentata. J. Neurophysiol. **79**, 1518–1534 (1998).
- 147. Buckmaster, P. S., Strowbridge, B. W. & Schwartzkroin, P. A. A comparison of rat hippocampal mossy cells and CA3c pyramidal cells. *J. Neurophysiol.* **70**, 1281–1299 (1993).
- Livsey, C. T. & Vicini, S. Slower spontaneous excitatory postsynaptic currents in spiny versus aspiny hilar neurons. *Neuron* 8, 745–755 (1992).
- 149. Strowbridge, B. W., Buckmaster, P. S. & Schwartzkroin, P. A. Potentiation of spontaneous synaptic activity in rat mossy cells. *Neurosci. Lett.* 142, 205–210 (1992).
- 150. Scharfman, H. E. & Schwartzkroin, P. A. Protection of dentate hilar cells from prolonged stimulation by intracellular calcium chelation. *Science* **246**, 257–260 (1989).
- 151. Scharfman, H. E. & Schwartzkroin, P. A. Responses of cells of the rat fascia dentata to prolonged stimulation of the perforant path: sensitivity of hilar cells and changes in granule cell excitability. *Neuroscience* 35, 491–504 (1990).
- 152. Li, G. *et al.* Hilar mossy cells share developmental influences with dentate granule neurons. *Dev. Neurosci.* **30**, 255–261 (2007).

Acknowledgements

This work is supported by the US National Institutes of Health, the Alzheimer's Association and the New York State Office of Mental Health.

Competing interests statement

The author declares no competing interests.